



12-1-05

IPW

Express Mail Label No. EV695506153US  
Date of Deposit: November 29, 2005

Attorney Docket No. 25371-021CIP

**IN THE UNITED STATES PATENT AND TRADEMARK OFFICE**

APPLICANT: Nakamura, et al.  
SERIAL NUMBER: 10/788,847 EXAMINER : Not Yet Assigned  
FILING DATE: February 27, 2004 ART UNIT : 1645  
FOR: GENE AND PROTEIN RELATING TO HEPATOCELLULAR CARCINOMA  
AND METHODS OF USE THEREOF

Commissioner for Patents  
P. O. Box 1450  
Alexandria, VA 22313-1450

**BEST AVAILABLE COPY**

**TRANSMITTAL**

Transmitted herewith for filing in the above-referenced patent application are the following documents:

1. Communication (1 pg);
2. Certified copy of Canadian Patent Application No. 2,399,569 (50 pgs);  
and
3. Return Postcard

Although Applicants believe that no additional fees are due in connection with this submission, the Commissioner is authorized to charge any deficiencies to Deposit Account No. 50-0311 (Reference No. 25371-021CIP).

If the enclosed papers are considered incomplete, the Mail Room is respectfully requested to contact the undersigned collect at (617) 542-6000, Boston, Massachusetts.

Respectfully submitted,

Ingrid A. Beattie, Reg. No. 42,306  
Cynthia A. Kozakiewicz, Reg. No. 42,764  
c/o MINTZ LEVIN  
Tel: (617) 542-6000  
Fax: (617) 542-2241  
Customer No. 30623

Dated: November 29, 2005



Express Mail Label No. EV695506153US  
Date of Deposit: November 29, 2005

Attorney Docket No. 25371-021CIP

**IN THE UNITED STATES PATENT AND TRADEMARK OFFICE**

APPLICANT: Nakamura, et al.  
SERIAL NUMBER: 10/788,847 EXAMINER : Not Yet Assigned  
FILING DATE: February 27, 2004 ART UNIT : 1645  
FOR: GENE AND PROTEIN RELATING TO HEPATOCELLULAR CARCINOMA  
AND METHODS OF USE THEREOF


Commissioner for Patents  
P. O. Box 1450  
Alexandria, VA 22313-1450

**COMMUNICATION**

Enclosed is a certified copy of Canadian Patent Application 2,399,569, the priority document of the above-referenced application.

The Commissioner is authorized to charge any fees that may be due, or to credit any overpayment, to the undersigned's account, Deposit Account No. 50-0311 Ref. No. 25371-021 CIP.

Respectfully submitted,

  
\_\_\_\_\_  
Ingrid A. Beattie, Reg. No. 42,306  
Cynthia A. Kozakiewicz, Reg. No. 42,764  
c/o MINTZ LEVIN  
Tel: (617) 542-6000  
Fax: (617) 542-2241  
Customer No. 30623

Dated: November 29, 2005



Office de la propriété  
intellectuelle  
du Canada

Un organisme  
d'Industrie Canada

Canadian  
Intellectual Property  
Office

An Agency of  
Industry Canada


*Bureau canadien  
des brevets  
Certification*

*Canadian Patent  
Office  
Certification*

La présente atteste que les documents  
ci-joints, dont la liste figure ci-dessous,  
sont des copies authentiques des docu-  
ments déposés au Bureau des brevets.

This is to certify that the documents  
attached hereto and identified below are  
true copies of the documents on file in  
the Patent Office.

Specification and Drawings, as originally filed, with Application for Patent Serial No:  
**2,399,569**, on August 23, 2002, by **YUSUKE NAKAMURA** and **YOICHI  
FURUKAWA**, for "Diagnostic Markers and Drug Targets for Treatment of Cancer".

  
Agent certificateur/Certifying Officer

November 23, 2005  
Date

Canada

(CIPO 68)  
31-03-04

OPIC  CIPO

## Abstract

To disclose mechanisms of hepatocellular carcinogenesis and to identify novel diagnostic markers and/or drug targets for treatment of hepatocellular carcinomas (HCCs), we analyzed expression profiles of clinical HCCs using a genome-wide cDNA microarray. From among the transcripts that were commonly up-regulated in these tumors, we identified a novel human gene, at chromosomal band 1p36.13, termed *DDEFL1* (development and differentiation enhancing factor-like 1) encoding a product that shared structural features with centaurin-family proteins as well as *VANGL1*, *LGN*, and a novel gene termed *ZNFN3A1* at chromosomal band 1q44. The deduced 903-amino-acid sequence of *DDFEL1* showed 46% homology to *DDEF/ASAP1* (development and differentiation enhancing factor), and contained an Arf GTPase-activating protein (ArfGAP) domain and two ankyrin repeats. The predicted 428-amino-acid sequence of *ZNFN3A1* contained a zf-MYND (MYND domain containing zinc finger protein) domain and a SET (Su (var) 3-9, Enhancer-of-zeste, Trithorax) domain. *VANGL1* (Van Gogh Like 1) is a gene homologous to *strabismus* (*Van Gogh*) that is involved in cell polarity and cell fate decisions in *Drosophila*. *LGN* protein interacts with alpha subunit of inhibitory heterotrimeric G proteinis ( $G\alpha_{i2}$ ). Gene transfer of *DDEFL1*, *ZNFN3A1* or *LGN* promoted proliferation of cells that lacked endogenous expression of either of these genes. Furthermore, reduction of *DDEFL1*, *ZNFN3A1*, *VANGL1* or *LGN* expression by transfection of their specific anti-sense S-oligonucleotides inhibited the growth of hepatocellular carcinoma cells. Our results provide novel insight into hepatocarcinogenesis and may contribute to development of new strategies for diagnosis and treatment of HCC.

## DIAGNOSTIC MARKERS AND DRUG TARGETS FOR TREATMENT OF CANCER

### Introduction

Hepatocellular carcinoma (HCC) is a leading cause of cancer deaths worldwide. In spite of recent progress in therapeutic strategies, prognosis of patients with advanced HCC remains very poor. Although molecular studies have revealed that alterations of *TP53*, *CTNNB1* and/or *AXIN1* genes can be involved in hepatocarcinogenesis (1, 2), these changes appear to be implicated in only a fraction of HCCs. To disclose mechanisms of hepatocellular carcinogenesis from a genome-wide point of view and to discover target molecules for diagnosis and for development of novel therapeutic drugs, we have been analyzing expression profiles of HCCs by means of a cDNA microarray representing 23,040 genes. These efforts have pinpointed 165 genes, including 69 ESTs, which appear to be up-regulated frequently in cancer tissues compared with corresponding non-cancerous liver cells. Since carcinogenesis involves activation of oncogenes and/or inactivation of tumor suppressor genes, enhanced expression of at least some of these 165 genes may reflect oncogenic properties.

In this paper we describe isolation of two novel human genes, *DDEFL1* (development and differentiation enhancing factor-like 1) and *ZNFN3A1* (Zinc Finger protein N3A1), and two human genes, *VANGL1* and *LGN* from among the transcripts whose expression was frequently elevated in HCCs. Reducing the expression of *DDEFL1*, *ZNFN3A1*, *VANGL1* or *LGN* by transfection of anti-sense DNA was able to inhibit growth of human HCC cells *in vitro*. These data should shed light on the mechanisms of hepatocellular carcinogenesis and also provide clues for development of novel diagnostic and therapeutic strategies.

## **Materials and Methods**

### **Patients and tissue specimens**

All HCC tissues and corresponding non-cancerous tissues were obtained with informed consent from surgical specimens of patients who underwent hepatectomy in Kyoto University Hospital.

### **Cell lines**

Human embryonic kidney 293 (HEK293) cells, COS7 cells, NIH3T3 cells and human hepatoma cell lines HepG2, Huh7 and Alexander were obtained from the American Type Culture Collection (ATCC). SNU423, SNU449, SNU475 were obtained from the Korea cell-line bank. All cells were grown in monolayers in appropriate media: Dulbecco's modified Eagle's medium for COS7, NIH3T3, HEK293, HepG2, Huh7 and Alexander and RPMI1640 for SNU423, SNU449 and SNU475; both media were supplemented with 10% fetal bovine serum and 1% antibiotic/antimycotic solution (Sigma).

### **RNA preparation and RT-PCR**

Total RNA was extracted with a Qiagen RNeasy kit (Qiagen) or Trizol reagent (Life Technologies, Inc.) according to the manufacturers' protocols. Ten-microgram aliquots of total RNA were reversely transcribed for single-stranded cDNAs using poly dT<sub>12-18</sub> primer (Amersham Pharmacia Biotech) with Superscript II reverse transcriptase (Life Technologies). Each single-stranded cDNA preparation was diluted for subsequent PCR amplification by standard RT-PCR experiments carried out in 20- $\mu$ l volumes of PCR buffer (TAKARA). Amplification proceeded for 4 min at 94°C for denaturing, followed by 20 (for *GAPDH*) or 33 (for *DDEFL1*) cycles of 94°C for 30 s, 56°C for 30 s, and 72°C for 45 s, in the GeneAmp PCR system 9700 (Perkin-Elmer, Foster City, CA). Primer sequences were; for *GAPDH*: forward, 5'-ACAACAGCCTCAAGA TCATCAG and reverse, 5'-GGTCCACCACTGACACGTTG; for *DDEFL1*: forward, 5'-AGCTGAGACATTTGTTCTCTTG and reverse: 5'-TATAAACCAG CTGAGTCCAGAG; for *ZNFN3A1* forward: 5'-TTCCCGATATCAACATC TACCAG reverse: 5'-AGTGTGTGACCTCAATAAGGCAT; for *VANGL1* forward: 5'-GAGTTGTATTATGAAGAGGCCGA reverse: 5'- ATGTCTCAGACTGTAA

GCGAAGG; and for *LGN* forward: 5'-ATCTGAAGCACTTAGCAATTGC reverse: 5'-CTGTAGCTCAGACCAAGAACC.

### **Northern-blot analysis**

Human multiple-tissue blots (Clontech, Palo Alto, CA) were hybridized with a <sup>32</sup>P-labeled cDNA designated *DDEFL1*, *ZNFN3A1* and *VANGL1*. Pre-hybridization, hybridization and washing were performed according to the supplier's recommendations. The blots were autoradiographed with intensifying screens at -80°C for 72 h.

### **5' rapid amplification of cDNA ends (5' RACE)**

5' RACE experiments were carried out using a Marathon cDNA amplification kit (Clontech, Palo Alto, CA) according to the manufacturer's instructions. For the amplification of the 5' part of *DDEFL1*, *ZNFN3A1*, and *VANGL1* cDNAs, gene-specific reverse primers (5'-CTCACTTGGCACGTCAG CAGGG, 5'-CTGCCAAGAAGTCGGAGTCTGGAG and 5'-TGTCAGCTCTCCG CTTGCGGAAAAAAG, respectively) and the AP-1 primer supplied in the kit were used. The cDNA template was synthesized from human liver mRNA. The PCR products were cloned using a TA cloning kit (Invitrogen) and their sequences were determined with an ABI PRISM 3700 DNA sequencer (Applied Biosystems).

### **Establishment of cells over-expressing *DDEFL*, *ZNFN3A1* or *LGN***

The entire-coding sequences of *DDEFL1*, *ZNFN3A1*, *LGN* or *LacZ* were cloned separately into the pcDNA3.1-myc/His, or pcDNA 3.1 vector (Invitrogen),. Each clone was transfected into NIH3T3 and SNU423 cells using FuGENE6 reagent (Boehringer) according to the supplier's recommendations. 24 h after transfection, geneticin was added to the cultures and single colonies were selected two weeks after transfection. Expression of *DDEFL1*, *ZNFN3A1*, and *LGN* was determined by semi-quantitative RT-PCR.

### **Immunoblotting**

Cells transfected with pcDNA3.1-myc/His-DDEFL1, pEGFP-ZNFN3A1, pFLAG-ZNFN3A1, pcDNA3.1-myc/His-VANGL1 or pcDNA3.1-myc/His-LGN were washed

twice with PBS and harvested in lysis buffer (150 mM NaCl, 1% Triton X-100, 50 mM Tris-HCl pH 7.4, 1mM DTT, and 1X complete Protease Inhibitor Cocktail (Boehringer)). After the cells were homogenized and centrifuged at 10,000xg for 30 min, the supernatant were standardized for protein concentration by the Bradford assay (Bio-Rad). Proteins were separated by 10% SDS-PAGE and immunoblotted with mouse anti-myc, anti-Flag, anti-GFP or anti-HA antibody. HRP-conjugated goat anti-mouse IgG (Amersham) served as the secondary antibody for the ECL Detection System (Amersham).

#### **Immunohistochemical staining**

Cells transfected with pcDNA3.1-myc/His-DDEFL1, pFLAG- ZNFN3A1 pcDNA3.1-myc/His-VANGL1 or pcDNA3.1-myc/His-LGN were fixed with PBS containing 4% paraformaldehyde for 15 min, then rendered permeable with PBS containing 0.1% Triton X-100 for 2.5 min at RT. Subsequently the cells were covered with 2% BSA in PBS for 24 h at 4°C to block non-specific hybridization. Mouse anti-myc monoclonal antibody (Sigma) at 1:1000 dilution or mouse anti-FLAG antibody (Sigma) at 1:2000 dilution was used for the first antibody, and the reaction was visualized after incubation with Rhodamine-conjugated anti-mouse second antibody (Leinco and ICN). Nuclei were counter-stained with 4',6'-diamidine-2'-phenylindole dihydrochloride (DAPI). Fluorescent images were obtained under an ECLIPSE E800 microscope.

#### **Colony-formation assay**

Cells plated onto 10-cm dishes ( $2 \times 10^5$  cells/dish) were transfected either with plasmid or with synthetic S-oligonucleotides, and cultured with an appropriate concentration of geneticin for two weeks. The cells were then fixed with 100% methanol and stained by Giemsa solution. Sense or antisense S-oligonucleotides of *DDEFL1*, encompassing the transcription-initiation site or the first ATG sequence, were transfected using LIPOFECTIN Reagent (GIBCO BRL). Sequences of the S-oligonucleotides were as follows: *DDEFL1* sense, 5'-CCTC GCGTGGTGAGTA; *DDEFL1* antisense, 5'-TACTCACCACGCGAGG, *ZNFN3A1* sense; 5'-GGCTCCATCCTCCCGC, *ZNFN3A1* antisense; 5'-GCGGGAGGATG GAGCC, *VANGL1* sense; 5'-ATACCCAATCCACTTA, *VANGL1* antisense; 5'-



TAAGTGGATTGGGTAT, *LGN* sense; 5'-CGATGGAGGAAAATTT, and *LGN* antisense; 5'-AAATTTTCCTCCATCG.

#### Flow cytometry

Cells were plated at a density of  $1 \times 10^5$  cells/100 mm dish and trypsinized at the given time course, followed by fixation in 70% cold ethanol. After RNase treatment, cells were stained with propidium iodide (50  $\mu$ g/ml) in PBS. Flow cytometry was performed on a Becton Dickinson FACScan and analyzed by CellQuest and ModFit software (Verity Software House). The percentages of nuclei in G0/G1, S and G2/M phases of the cell cycle, and any sub-G1 population were determined from at least 20,000 ungated cells.

#### 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) assay

Cells were plated at a density of  $5 \times 10^5$  cells/100 mm dish. At 24 hours after seeding, the cells were transfected in triplicate with sense or antisense S-oligonucleotide designated to suppress either *ZNFN3A1* or *VANGL1*. At 72 hours after transfection, the medium was replaced with fresh medium containing 500  $\mu$ g/ml of MTT (3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyl tetrazolium bromide) (Sigma) and the plates were incubated for four hours at 37°C. Subsequently, the cells were lysed by the addition of 1 ml of 0.01 N HCl/10%SDS and absorbance of lysates was measured with an ELISA plate reader at a test wavelength of 570 nm (reference, 630 nm). The cell viability was represented by the absorbance compared to that of control cells.

## Results

### Identification of *DDEFL1*, a commonly up-regulated gene in human hepatocellular carcinomas

By means of a genome-wide cDNA microarray containing 23040 genes, we compared expression profiles of 20 HCCs with their corresponding non-cancerous liver tissues. A gene with an in-house accession number of B9362 corresponding to an EST, Hs.44579 of a UniGene cluster (<http://www.ncbi.nlm.nih.gov/UniGene/>), was found to be over-expressed in a range between 1.57 and 5.83 (Fig. 1a). Since an open reading frame of this gene encoded a protein approximately 60% identical

to that of DDEF2\_(development and differentiation enhancing factor\_2), we termed this gene *DDEFL1* (development and differentiation enhancing factor-like 1). To clarify the results of the cDNA microarray, we examined expression of this transcript in an additional 11 HCCs by semi-quantitative RT-PCR and confirmed its increased expression in nine of these tumors (Fig. 1b).

### **Isolation and structure of a novel gene *DDEFL1***

Multiple-tissue northern-blot analysis using a PCR product of *DDEFL1* as a probe revealed a 4-kb transcript that was expressed in lung, liver, small intestine, placenta and peripheral blood leukocyte (Fig. 2a). Since *B9362* was smaller than that detected on the Northern blot, we carried out 5'RACE experiments to determine the entire coding sequence of the gene. The complete cDNA consisted of 4050 nucleotides, with an open reading frame of 2712 nucleotides encoding a 903-amino-acid protein (GenBank accession number AB051853). The first ATG was flanked by a sequence (CCCGCCATGC) that agreed with the consensus sequence for initiation of translation in eukaryotes, with an in-frame stop codon upstream. Using the BLAST program to search for homologies in the NCBI (the National Center for Biotechnology Information) database, we identified a genomic sequence with GenBank accession number AL357134, which had been assigned to chromosomal band 1p36.12. Comparison of the cDNA and genomic sequences disclosed that the novel gene consisted of 25 exons (Fig. 2b).

A search for protein motifs with the Simple Modular Architecture Research Tool (SMART, <http://smart.embl-heidelberg.de>) revealed that the predicted protein contained two coiled-coil regions (codons 141-172 and 241-278), a PH (Pleckstrin homology) motif (codons 303-396), a motif of ArfGAP (GTPase-activating protein for Arf) (codons 426-551) and two ankyrin repeats (codons 585-617 and 621-653). This structure was similar to centaurin beta 1 and centaurin beta 2 (Fig. 2c). In particular, DDEFL1 shared features of centaurin-family proteins such as a PH domain, a target of phosphatidylinositol 3,4,5-trisphosphate, and a motif of ArfGAP. The amino acid sequence of the ArfGAP motif of DDEFL1 was 67.8% identical to that of DDEF2 (Fig. 2d). Notably, the CXXCX<sub>16</sub>CXXC motif, representing a zinc finger structure essential to GAP activity, was completely preserved.

### **Subcellular localization of DDEFL1**

We transiently transfected a plasmid expressing myc-tagged DDEFL1 protein (pDNA-myc/His-DDEFL1) into COS7 cells and subjected the cells to immunocytochemical staining. The expected myc-tagged protein was detected on western blots with an anti-myc antibody (Fig. 3a), and subsequent microscopic analysis indicated that the protein was present mainly in the cytoplasm (Fig. 3b). DDEFL1 was also localized in the cytoplasm of human embryonal kidney (HEK293) cells.

### **Effect of *DDEFL1* on cell growth**

We carried out a colony-formation assay by transfecting NIH3T3 cells with a plasmid expressing *DDEFL1* (pcDNA-DDEFL1). Cells transfected with pcDNA-DDEFL1 produced markedly more colonies than control cells. An increase in colony formation similarly occurred with transfected human hepatoma SNU423 and Alexander cells, in which endogenous expression of *DDEFL1* is very low (Fig. 4a).

To investigate this growth-promoting effect further, we established NIH3T3 cells that stably expressed exogenous *DDEFL1* (Fig. 4b). The growth rate of NIH3T3-DDEFL1 cells was significantly higher than that of parental NIH3T3 cells in culture media containing 10% FBS (Fig. 4c); but although NIH3T3-DDEFL1 cells survived for 6 days in media containing only 0.1% FBS, control NIH3T3 cells died within 6 days under the same conditions (Fig. 4d).

### **Suppression of *DDEFL1* expression in human hepatoma SNU475 cells by antisense S-oligonucleotides**

Six pairs of control and antisense S-oligonucleotides corresponding to the *DDEFL1* gene were transfected into SNU475 cells, which had shown the highest level of *DDEFL1* expression among the six hepatoma cell lines we examined. Antisense S-oligonucleotides #1 and #5 significantly suppressed expression of *DDEFL1* compared to control S-oligonucleotides, 12 and 24 hours after transfection (Fig. 5a). Six days after transfection, surviving cells transfected with antisense S-oligonucleotides #1 or #5 were markedly fewer than cells transfected with control S-oligonucleotides (Fig. 5b). Consistent results were obtained in three independent experiments.

**Identification of A6681, a novel gene frequently up-regulated in HCCs**

Among commonly up-regulated genes by the microarray analysis, we also focused on a gene, A6681, corresponding to an EST (Hs.8109) of a UniGene cluster (<http://www.ncbi.nlm.nih.gov/UniGene/>), because its expression was significantly up-regulated in eleven of twelve clinical HCCs compared with the corresponding non-cancerous liver tissues (Fig. 6a). The elevated expression of A6681 was also confirmed in additional ten HCC cases by semi-quantitative RT-PCR (Fig. 6b).

**Expression, isolation and characterization of a novel human gene, ZNFN3A1**

Multi-tissue northern blot analysis using A6681 cDNA as a probe showed a 1.7-kb transcript expressed specifically in testis and skeletal muscle (Fig. 7a). We searched for genomic sequences corresponding to A6681 in the genomic databases and found a draft sequence assigned to chromosomal band 1q44 (GenBank accession number:NT004536). Using GENSCAN and Gene Recognition and Assembly Internet Link program, we obtained candidate-exon sequences and performed exon-connection using these sequence information. In addition, we carried out 5'-RACE to determine the sequence of the 5' region of the transcript. As a result, we obtained an assembled sequence of 1622 nucleotides containing an open reading frame of 1284 nucleotides (GenBank accession number:AB057595). Simple Modular Architecture Research Tool suggested that the predicted protein contained a zf-MYND [zinc finger protein (MYND domain containing)] domain (codons 49-87) and a SET [(Su (var) 3-9, Enhancer-of-zeste, Trithorax)] domain (codons 117-246) (Fig. 7b). The gene was termed *ZNFN3A1* (zinc finger protein, subfamily 3A (MYND domain containing), 1) by the nomenclature committee.

**Subcellular localization of ZNFN3A1**

The entire coding region corresponding to *ZNFN3A1* was cloned into a pEGFP-N1 vector and a pFlag-CMV-5a vector, and these constructs were transiently transfected into SNU475 cells. ZNFN3A1-EGFP fusion protein was present in the cytoplasm and nucleus by fluorescent microscopy. Similarly flag-tagged ZNFN3A1

protein was detected in the cytoplasm and nucleus by fluorescent immunohistochemical staining (Fig. 8).

#### **Effect of *ZNFN3A1* on cell growth**

To analyze the effect of *ZNFN3A1* on cell growth, we carried out a colony-formation assay by transfecting NIH3T3 cells with a plasmid expressing *ZNFN3A1* (pcDNA-*ZNFN3A1*). Compared with mock and antisense-*ZNFN3A1* expressing plasmids, sense-*ZNFN3A1* expressing plasmid induced markedly more colonies in NIH3T3 cells (Fig. 9a, b). This result was confirmed by three independent experiments. To investigate this growth-promotive effect of *ZNFN3A1*, we established NIH3T3 cells that stably expressed exogenous *ZNFN3A1* (NIH3T3-*ZNFN3A1* cells) (Fig. 9c). As shown Fig. 9d, growth rate of NIH3T3-*ZNFN3A1* cells was higher than those transfected with antisense *ZNFN3A1* or mock cells.

#### **Growth suppression of hepatoma cells by antisense S-oligonucleotides designated to reduce expression of *ZNFN3A1***

To test whether suppression of *ZNFN3A1* may result in cell cycle arrest and/or cell death of HCC cells, we synthesized various antisense S-oligonucleotides designated to suppress its expression. Among them, antisense S-oligonucleotides (AS) encompassing the initiation codon, but not other antisense or control S-oligonucleotides (SE) significantly decreased endogenous expression of *ZNFN3A1* in SNU475 and Huh7 cells that constitutively express *ZNFN3A1* abundantly (Fig. 10a). Transfection of the antisense S-oligonucleotides (AS) significantly reduced number of surviving cells compared with that of control sense S-oligonucleotides (SE) (Fig. 10b, c). This result was confirmed by three independent experiments. Furthermore, FACS analysis demonstrated that inhibition of *ZNFN3A1* significantly decreased number of cells in S phase and increased number in sub-G1 phase (Fig. 10d). These results suggest that *ZNFN3A1* may play an important role for cell growth and/or survival of hepatocellular carcinoma cells.

#### ***D3244* is commonly increased in human hepatocellular carcinomas**

Additionally, we focused on a gene, *D3244* corresponding to an EST

(Hs.122730) of a UniGene cluster (<http://www.ncbi.nlm.nih.gov/UniGene/>), because it was also significantly up-regulated in ten of twelve clinical HCCs compared with the corresponding non-cancerous liver tissues. The relative expression ratio compared to corresponding non-cancerous tissue of these 12 tumors ranged from 1.5 to 16.0 (Fig. 11a). The elevated expression of *D3244* was also confirmed in ten additional HCC cases by semi-quantitative RT-PCR (Fig. 11b).

### **Expression of *VANGL1* in human adult tissues**

Multi-tissue northern blot analysis using *D3244* cDNA as a probe showed a 1.9-kb transcript abundantly expressed in testis and ovary in a tissue-specific manner (Fig. 12a). We searched for genomic sequences corresponding to *D3244* in databases in NCBI (<http://www.ncbi.nlm.nih.gov/>), and found two sequences (GenBank accession number:AL450389 and AL592436) assigned to chromosomal band 1p22. Using GENSCAN, and Gene Recognition and Assembly Internet Link program, we predicted candidate-exon sequences and performed exon-connection. In addition, we carried out 5' RACE to determine the sequence of the 5' region of the transcript. As a result, we obtained an assembled human cDNA sequence of 1879 nucleotides containing an open reading frame of 1572 nucleotides (GenBank accession number: AB057596). The predicted amino acid sequence shared 40% and 63% identity with *strabismus* (*Van Gogh*) and *VANGL2*. Hence, we termed the gene corresponding *D3244* as *VANGL1* (*Van Gogh Like 1*). Simple Modular Architecture Research Tool suggested that the predicted protein contained putative four transmembrane domains (codons 111-133,148-170,182-204, 219-241) (Fig. 12b).

### **Subcellular localization of *VANGL1***

The pcDNA3.1-myc/His-*VANGL1* plasmid expressing c-myc-tagged *VANGL1* protein was transiently transfected into SNU475 cells. Immunocytochemical staining revealed that the tagged *VANGL1* protein was present in the cytoplasm (Fig. 13).

### **Growth suppression of hepatoma cells by antisense S-oligonucleotides designated to reduce expression of *VANGL1***

To test whether suppression of *VANGL1* may result in cell cycle arrest and/or cell death of HCC cells, we synthesized antisense S-oligonucleotides to suppress its expression. Among them, antisense S-oligonucleotides encompassing the initiation codon significantly decreased endogenous expression of *VANGL1* in SNU475 cells (Fig. 14a). Transfection of the antisense S-oligonucleotides significantly reduced number of surviving cells compared with control sense S-oligonucleotides (Fig. 14b, c). This result was confirmed by three independent experiments. Furthermore, FACS analysis demonstrated that inhibition of *VANGL1* significantly increased number of cells at sub-G1 phase (Fig. 14d). These results suggest that *VANGL1* may play an important role for cell growth and/or survival of hepatocellular carcinoma cells.

#### ***LGN* is commonly increased in human hepatocellular carcinomas**

Among commonly up-regulated genes by the microarray analysis, we further selected a gene, *D3636* corresponding to *LGN* (GenBank accession number: U54999), because it was significantly up-regulated in ten of twelve clinical HCCs compared with the corresponding non-cancerous liver tissues. The relative expression ratio compared to corresponding non-cancerous tissue of these 12 tumors ranged from 0.7 to 16.0 (Fig. 15a). The elevated expression of *LGN* was also confirmed in additional ten HCC cases by semi-quantitative RT-PCR (Fig. 15b).

#### **Genomic structure of *LGN***

*LGN* cDNA consists of 2,336 nucleotides and encodes a 677 amino acid peptide. Comparison of the cDNA sequence with genomic sequences disclosed that the *LGN* gene consists of 14 exons (Fig. 16).

#### **Subcellular localization of *LGN***

The pcDNA3.1-myc/His-*LGN* plasmid expressing c-myc-tagged *LGN* protein was transiently transfected into COS7 cells. A 72 kDa-band corresponding to myc-tagged *LGN* protein was detected by immunoblot analysis (Fig. 17). Immunocytochemical staining revealed that the tagged *LGN* protein was present in the cytoplasm and nucleus in the cells.

**LGN gene transfer can promote cell growth**

To analyze the effect of *LGN* on cell growth, we carried out a colony-formation assay by transfecting NIH3T3, SNU423, Alexander and SNU475 cells with a plasmid expressing *LGN* (pcDNA3.1-myc/His-*LGN*). Compared with a control plasmid (pcDNA3.1-myc/His-LacZ), pcDNA3.1-myc/His-*LGN* produced markedly a larger number of colonies in these cells (Fig. 18a). This result was confirmed by three independent experiments.

To further investigate the effect of *LGN* on cell growth, we established NIH3T3 cells that stably expressed exogenous *LGN* (NIH3T3-*LGN* cells). NIH3T3-*LGN* cells showed higher growth rate than control NIH3T3-*LacZ* cells (Fig. 18b).

**Antisense S-oligonucleotides of *LGN* suppressed growth of human hepatoma SNU475 cells**

We synthesized five pairs of sense and antisense S-oligonucleotides corresponding to *LGN* and then transfected into SNU423 cells. When we transfected antisense S-oligonucleotides, the antisense S-oligonucleotides #4 significantly suppressed expression of *LGN* compared to control S-oligonucleotides, 12 hours after transfection (Fig. 19a). Six days after transfection, the number of surviving cells transfected with antisense S-oligonucleotides #4 were markedly fewer than that with control S-oligonucleotides (Fig. 19b). Consistent results were obtained in three independent experiments.

**Discussion**

cDNA microarray technologies have enabled us to obtain comprehensive profiles of gene expression in normal versus malignant cells (1, 3-4). This approach discloses the complex nature of cancer cells, and helps to improve our understanding of carcinogenesis. Identification of genes that are deregulated in tumors can lead to more precise and accurate diagnosis of individual cancers, and to development of novel therapeutic targets (7). The genome-wide analysis of HCCs we reported previously disclosed a number of genes associated with



carcinogenesis, as well as genes affecting differentiation and vessel invasion. To identify ideal therapeutic targets for treatment of HCCs, we focused in the present work on *DDEFL1*, one of the genes up-regulated most frequently in HCCs according to our microarray analysis.

Since DDEFL1 showed 46% identity to a member of the centaurin family, DDEF2 (development and differentiation enhancing factor 2), a protein that regulates re-organization of the actin cytoskeleton, DDEFL1 may also play a role in organization of cellular structure (8). Because DDEFL1 also conserves a PH domain and an ArfGAP motif it appears to be a new member of the centaurin family, regulating Arf small GTPase by means of GAP activity. The PH domain, observed in the majority of molecules belonging to the Dbl family of GEFs, is thought to play a crucial role in relocation of proteins by interacting with specific target molecules and/or by directly regulating catalytic domains (5,9). Although DDEF2 is localized in peripheral focal adhesions, we found myc-tagged DDEFL1 protein to be diffuse in cytoplasm. Hence DDEFL1 may have a function different from DDEF2, but further investigation using specific antibody will be required to disclose the sub-cellular location(s) of DDEFL1 more precisely.

Arf proteins have been implicated in important cellular processes such as vesicular membrane transport, maintenance of the integrity of ER and Golgi compartments, and regulation of the peripheral cytoskeleton (6). Six members of Arf family (Arf1-Arf6) and their functions have been identified so far (11). For example, Arf6 proteins have been implicated as regulators of the cytoskeleton to alter the morphology of focal adhesions and to block spreading of cells, and DDEF2 displays GAP activity toward Arf1. To disclose the function of DDEFL1, however, it will be necessary to identify the molecular targets of its ArfGAP domain.

Since in our experiments over-expression of *DDEFL1* promoted growth promotion and survival of cells under low-serum conditions, *DDEFL1* may provide a growth advantage to cancer cells in poor nutritional and hypoxic conditions. Although the mechanism of this mitogenic effect remains to be investigated, the frequent up-regulation of *DDEFL1* in HCCs underscores the importance of this gene in hepatocarcinogenesis.

ZNFN3A1 has a zf-MYND domain and a SET domain, both of which are conserved in a putative proto-oncogene, *MTG8/ETO*. *MTG8/ETO* was identified by

molecular characterization of a t(8;21)(q22;q22) translocation breakpoint commonly found in the subset M2 of acute myeloid leukemias (12). The chimeric protein AML1/MTG8 produced by the translocation may play a crucial role in leukemogenesis by inactivating the function of AML1 and/or activating transcription through the zf-MYND domain of MTG8 (13, 14). Therefore up-regulation of *ZNFN3A1* may promote cell proliferation by regulating transcription of various genes as a component of transcriptional complex.

*Strabismus* (*stbm*) was identified as a gene responsible for a mutant fruit fly with rough eye phenotype (15). The gene is required to maintain polarity in the eye, legs and bristles and to decide cell fate of R3 and R4 photoreceptors in the *Drosophila*. A mouse gene homologous to *stbm*, *Ltap*, was altered in the neural tube mutant mouse Loop-tail, which is a human model of neural tube defects (NTDs) (16). Hence, *VANGL1* may also play important roles in cellular polarity, cell fate decision, and/or organization of tissues. Since *VANGL1* is frequently up-regulated in HCCs and suppression of its expression significantly reduced growth or survival of cancer cells, *VANGL1* may confer prolonged survival and/or depolarized growth to cancer cells.

*LGN* protein was previously reported as a protein interacting with alpha subunit of inhibitory heterotrimeric G proteins ( $G\alpha_{i2}$ ) (17). The activating mutations of  $G\alpha_{i2}$  have ever been reported in pituitary tumor and other endocrine tumors (18, 19, 20). However, involvement of *LGN* in tumorigenesis or carcinogenesis has not yet been reported. In this study, we showed that *LGN* was frequently up-regulated in HCCs compared with non-cancerous liver tissues. In addition, colony formation assay suggested that *LGN* might have oncogenic activity. Taken together, enhanced expression of *LGN* may activate  $G\alpha_{i2}$  and mediate oncogenic signals in hepatocarcinogenesis. Further analysis of function of *LGN* and signal transduction pathway of heterotrimeric G proteins ( $G\alpha_{i2}$ ) is required for a more profound understanding of hepatocarcinogenesis.

We have demonstrated here that suppressing the expression of *DDEFL1*, *ZNFN3A1*, *VANGL1* or *LGN* by antisense oligonucleotides markedly decreases growth of HCC cells. Although the precise molecular mechanism by which the antisense S-oligonucleotides can suppress growth needs to be clarified, our data clearly indicate that these genes could be good candidates as diagnostic markers

for liver cancers, and could represent molecular targets for development of effective drugs to treat this often intractable disease.

## References

1. Perou, C. M., Sorlie, T., Eisen, M. B., van de Rijn, M., Jeffrey, S. S., Rees, C. A., Pollack, J. R., Ross, D. T., Johnsen, H., Akslen, L. A., Fluge, O., Pergamenschikov, A., Williams, C., Zhu, S. X., Lonning, P. E., Borresen-Dale, A. L., Brown, P. O., and Botstein, D. Molecular portraits of human breast tumours, *Nature*. **406**: 747-752, 2000.
2. Satoh, S., Daigo, Y., Furukawa, Y., Kato, T., Miwa, N., Nishiwaki, T., Kawasoe, T., Ishiguro, H., Fujita, M., Tokino, T., Sasaki, Y., Imaoka, S., Murata, M., Shimano, T., Yamaoka, Y., and Nakamura, Y. AXIN1 mutations in hepatocellular carcinomas, and growth suppression in cancer cells by virus-mediated transfer of AXIN1, *Nat Genet*. **24**: 245-250, 2000.
3. Clark, E. A., Golub, T. R., Lander, E. S., and Hynes, R. O. Genomic analysis of metastasis reveals an essential role for RhoC, *Nature*. **406**: 532-535, 2000.
4. Okabe, H., Satoh, S., Kato, T., Kitahara, O., Yanagawa, R., Yamaoka, Y., Tsunoda, T., Furukawa, Y., and Nakamura, Y. Genome-wide analysis of gene expression in human hepatocellular carcinomas using cDNA microarray: identification of genes involved in viral carcinogenesis and tumor progression, *Cancer Res*. **61**: 2129-2137, 2001.
5. Jackson, T. R., Kearns, B. G., and Theibert, A. B. Cytohesins and centaurins: mediators of PI 3-kinase-regulated Arf signaling, *Trends Biochem Sci*. **25**: 489-495, 2000.
6. Cukierman, E., Huber, I., Rotman, M., and Cassel, D. The ARF1 GTPase-activating protein: zinc finger motif and Golgi complex localization, *Science*. **270**: 1999-2002, 1995.
7. Golub, T. R., Slonim, D. K., Tamayo, P., Huard, C., Gaasenbeek, M., Mesirov, J. P., Coller, H., Loh, M. L., Downing, J. R., Caligiuri, M. A., Bloomfield, C. D., and Lander, E. S. Molecular classification of cancer: class discovery and class prediction by gene expression monitoring, *Science*. **286**: 531-537, 1999.
8. Randazzo, P. A., Andrade, J., Miura, K., Brown, M. T., Long, Y. Q., Stauffer, S., Roller, P., and Cooper, J. A. The Arf GTPase-activating protein ASAP1 regulates the actin cytoskeleton, *Proc Natl Acad Sci U S A*. **97**: 4011-4016, 2000.
9. Cerione, R. A., and Zheng, Y. *Curr. Opin. Cell. Biol.* **8**: 216-222, 1996.

- Chardin, P., Paris, S., Antonny, B., Robineau, S., Beraud-Dufour, S., Jackson, C. L., and Chabre, M. A human exchange factor for ARF contains Sec7- and pleckstrin-homology domains. *Nature*. 384: 481-484, 1996.
10. Ren, R., Mayer, B. J., Cicchetti, P., and Baltimore, D. Identification of a ten-amino acid proline-rich SH3 binding site. *Science*. 259: 1157-1161, 1993.
  11. Moss, J., Vaughan, M. Structure and function of ARF proteins: activators of cholera toxin and critical components of intracellular vesicular transport processes. *J Biol Chem*. 270: 12327-12330, 1995.
  12. Frank RC, Sun X, Berguido FJ, Jakubowiak A, Nimer SD. The t(8;21) fusion protein, AML1/ETO, transforms NIH3T3 cells and activates AP-1. *Oncogene*. 18: 1701-1710, 1999 .
  13. Gelmetti V, Zhang J, Fanelli M, Minucci S, Pelicci PG, Lazar MA. Aberrant recruitment of the nuclear receptor corepressor-histone deacetylase complex by the acute myeloid leukemia fusion partner ETO. *Mol Cell Biol*. 18: 7185-7191, 1998.
  14. Anderson SF, Schlegel BP, Nakajima T, Wolpin ES, Parvin JD. BRCA1 protein is linked to the RNA polymerase II holoenzyme complex via RNA helicase A. *Nat Genet*. 19: 254-256, 1998.
  15. Wolff T, Rubin GM. Strabismus, a novel gene that regulates tissue polarity and cell fate decisions in *Drosophila*. *Development* 125:1149-1159, 1998.
  16. Kibar Z, Vogan KJ, Groulx N, Justice MJ, Underhill DA, Gros P. Ltap, a mammalian homolog of *Drosophila* Strabismus/Van Gogh, is altered in the mouse neural tube mutant Loop-tail. *Nat Genet*. 28: 251-255, 2001.
  17. Mochizuki, N., Cho, G., Wen, B., Insel, P. A. Identification and cDNA cloning of a novel human mosaic protein, LGN, based on interaction with G<sub>α2</sub>. *Gene*. 181: 39-43, 1996.
  18. Hermouet, S., Merendino, J. J. Jr., Gutkind, J. S., Spiegel, A. M. Activating and inactivating mutations of the alpha subunit of Gi2 protein have opposite effects on proliferation of NIH 3T3 cells. *Proc Natl Acad Sci USA*. 88: 10455-10459, 1991.
  20. Pace, A. M., Wong, Y. H., Bourne, H. R. A mutant alpha subunit of Gi2 induces neoplastic transformation of Rat-1 cells. *Proc Natl Acad Sci USA*. 88: 7031-7035, 1991.205.

21. Lyons, J., Landis, C. A., Harsh, G., Vallar, L., Grunewald, K., Feichtinger, H., Duh, Q. Y., Clark, O. H., Kawasaki, E., Bourne, H. R., et al. Two G protein oncogenes in human endocrine tumors. *Science*. 249: 655-659, 1990.

### Figure Legends

Figure 1. Expression of *B9362* in HCCs a) Relative expression ratio (cancer/non-cancer) of *B9362* in primary 20 HCCs examined by cDNA microarray. Its up-regulated expression (Cy3:Cy5 intensity ratio, >2.0) was observed in 11 of the 12 HCCs that passed through the cutoff filter (both Cy3 and Cy5 signals greater than 25,000). b) Expression of *B9362* was analyzed by semi-quantitative RT-PCR using additional 11 HCC cases. Expression of *GAPDH* served as an internal control.

Figure 2. a) Northern blot analysis of *DDEFL1* in various human tissues. The transcript of *B9362* is approximately 4-kb by size. b) The structure of *DDEFL1*. The *DDEFL1* cDNA consists of 4,050 nucleotides with an ORF of 2,712 nucleotides and is composed of 25 exons. c) Similarity between the expected *DDEFL1* protein and members of ArfGAP family. d) Identity between the amino acid sequence of the ArfGAP motif in *DDEFL1* and that in *DDEF2*. The arrows indicate a CXXCX<sub>16</sub>CXXC motif, representing a zinc finger structure essential to GAP activity.

Figure 3. Subcellular localization of *DDEFL1*. a) The western blot analysis showed that cMyc-tagged *DDEFL1* protein was expressed in COS7 cells transfected with pcDNA-*DDEFL1*-myc plasmid. b) Immunocytochemistry of the cells suggested that cMyc-tagged *DDEFL1* protein localized in the cytoplasm.

Figure 4. Growth-promoting effect of *DDEFL1*. a) Colony formation assays show that *DDEFL1* promotes cell growth in NIH3T3, SNU423, and Alexander cells. b) NIH3T3-*DDEFL1* cells stably expressed exogenous *DDEFL1*. c) Growth of NIH3T3-*DDEFL1* cells stably expressing exogenous *DDEFL1* was statistically higher than that of mock (NIH3T3-LacZ) cells in culture media containing 10% FBS ( $P < 0.05$ ). d) Growth of NIH3T3-*DDEFL1* cells was statistically higher than that of mock cells in culture media containing 0.1% FBS ( $P < 0.01$ ).

Figure 5. Growth suppression by antisense S-oligonucleotides designated to suppress *DDEFL1* in SNU475 cells. a) Designation of antisense S-oligonucleotides. Reduced expression of *DDEFL1* by the transfection of AS1 or AS5 antisense S-oligonucleotides. b) AS1 and AS5 suppressed growth of SNU475 cells.

Figure 6. Expression of *A6681* in HCCs a) Relative expression ratios (cancer/non-cancer) of *A6681* in primary 20 HCCs examined by cDNA microarray. Its up-regulated expression (Cy3: Cy5 intensity ratio, >2.0) was observed in 11 of the 12 HCCs that passed through the cutoff filter (both Cy3 and Cy5 signals greater than 25,000). B) Expression of *A6681* was analyzed by semi-quantitative RT-PCR using additional 10 HCC cases. T, tumor tissue; N, normal tissue. Expression of *GAPDH* served as an internal control.

Figure 7. a) Multiple-tissue northern blot analysis of *ZNFN3A1* in various human tissues. b) Predicted protein structure of *ZNFN3A1*.

Figure 8. Subcellular localization of *ZNFN3A1* a) SNU475 cells were transfected with pEGFP- *ZNFN3A1*. Nuclei were counter-stained with DAPI. b) SNU475 cells transfected with pFLAG-*ZNFN3A1* were stained with anti-Flag monoclonal antibody and visualized by Rhodamine conjugated secondary anti-mouse IgG antibody. Nuclei were counter-stained with DAPI.

Figure 9. Effect of *ZNFN3A1* on cell growth. a) Colony formation assay of *ZNFN3A1* in NIH3T3 cells. b) Number of colonies counted by electric densitometry. c) Expression of *ZNFN3A1* in mock, NIH3T3-antisense *ZNFN3A1*, and NIH3T3-*ZNFN3A1* cells d) Cell growth of mock, NIH3T3-antisense *ZNFN3A1*, and NIH3T3-*ZNFN3A1* cells.

Figure 10 Growth suppressive effect of antisense S-oligonucleotides designated to suppress *ZNFN3A1*. a) Expression of *ZNFN3A1* in SNU475 cells treated with either control (SE) or antisense (AS) oligonucleotides for 12 h. b) S-

oligonucleotides (AS) suppressed growth of SNU423 cells. c) Analysis of cell viability by MTT assay. d) FACS analysis of Huh7 cells treated with control (SE) or antisense (AS) oligonucleotides.

**Figure 11. Expression of *VANGL1* in HCCs** a) Relative expression ratios (cancer/non-cancer) of *VANGL1* in primary 20 HCCs examined by cDNA microarray. Its up-regulated expression (Cy3:Cy5 intensity ratio, >2.0) was observed in 10 of the 12 HCCs that passed through the cutoff filter (both Cy3 and Cy5 signals greater than 25,000). b) Expression of *A6681* was analyzed by semi-quantitative RT-PCR using additional 10 HCC cases. T, tumor tissue; N, normal tissue. Expression of *GAPDH* served as an internal control.

**Figure 12.** a) Multiple-tissue northern blot analysis of *VANGL1* in various human tissues. b) Predicted protein structure of *VANGL1*.

**Figure 13. Subcellular localization of *VANGL1*** a) SNU475 cells transfected with pcDNA3.1-myc/His-LGN were stained with mouse anti-myc monoclonal antibody and visualized by Rhodamine conjugated secondary anti-mouse IgG antibody. Nuclei were counter-stained with DAPI.

**Figure 14** Growth suppressive effect of antisense S-oligonucleotides designated to suppress *VANGL1*. a) Expression of *VANGL1* in SNU475 cells treated with either control or antisense oligonucleotides for 12 h. b) S-oligonucleotides suppressed growth of SNU423 cells. c) Analysis of cell viability by MTT assay. d) FACS analysis of cells treated with sense or antisense oligonucleotides.

**Figure 15** LGN gene expression of HCCs compared with their corresponding non-cancerous liver tissues. a) Relative expression ratios (cancer/non-cancer) of *LGN* in primary 20 HCCs studied by cDNA microarray. Its up-regulated expression (Cy3:Cy5 intensity ratio, >2.0) was observed in 10 of the 12 HCCs that passed through the cutoff filter (both Cy3 and Cy5 signals greater than 25,000). b) Expression of *LGN* was analyzed by semi-quantitative RT-PCR using additional ten HCCs. RT-PCR experiments were performed with a *LGN*-specific primer set using



RNA extracted from HCCs. Expression of *GAPDH* served as an internal control. T, tumor tissue; N, normal tissue.

Figure 16. Genomic structure of *LGN*.

Figure 17. Subcellular localization of *LGN*. COS7 cells transfected with pcDNA3.1-myc/His-*LGN* were stained with mouse anti c-myc antibody and visualized by Rhodamine conjugated secondary anti-mouse IgG antibody. Nuclei were counter-stained with DAPI.

Figure 18. Growth-promoting effect of *LGN*. a) Colony formation assays show that *LGN* promotes cell growth in NIH3T3, SNU423, Alexander, and SNU423 cells. b) Growth of NIH3T3-*LGN* cells stably expressing exogenous *LGN* was higher than that of mock (NIH3T3-LacZ) cells in culture media containing 10% FBS.

Figure 19. Growth suppression by antisense S-oligonucleotides designated to suppress *LGN* expression in human hepatoma SNU423 cells. a) Reduced expression of *LGN* by the transfection of antisense S-oligonucleotides #4. b) S-oligonucleotides #4 suppressed growth of SNU423 cells.

## Sequence\_Table

SEQ ID 1

DDEF1(nucleotide)

GTGCCCCCGCGCTCCGCTCCGGCAGCTCCACGCTCGCGCCCGCCATGCCGGAGCA  
GTTTCAGCGTCGCCGAGTTCCTGGCCGTACCCGCGGAGGACCTCAGCTCCCCGGCTG  
GGGCCGCCGCTTCGCCGCCAAGATGCCCGGTACCGAGGGGCGGCGCTGGCGCG  
GGAGGAGATCTTGGAAGGAGACCAAGCCATCCTGCAGAGAATAAAGAAGGCTGTGCG  
GGCAATCCATAGCTCCGGCCTTGGCCATGTGGAGAATGAAGAGCAGTACCGAGAGGC  
CGTGGAATCCTTAGGCAACAGCCACCTGTCCCAAGAACAGCCATGAGCTGTCCACAGG  
CTTCCTAAACTTGGCCGTGTTACCCGCGAGGTTGCTGCGCTCTTCAAGAACCTGATT  
CAGAACTTGAACAACATTGTCTCTTCCCCCTGGACAGTCTGATGAAGGGGCAGCTGA  
GGGACGGTCGACAGGATTCCAAAAAACAGCTGGAGAAGGCATGGAAGGACTATGAAG  
CCAAATGGCCAAGCTGGAGAAGGAGCGCGATCGGGCCAGGGTGACAGGAGGGATC  
CCTGGGGAGGTGGCCCAGGACATGCAGAGAGAGCGGCGCATCTTCCAGCTGCACAT  
GTGTGAGTATCTGCTCAAAGCCGGGGAGAGCCAGATGAAGCAAGGTCCTGACTTCCTT  
CAGAGCCTCATCAAGTTCCTCCACGCCCAGCACAACCTTTTCCAAGATGGCTGGAAGG  
CTGCCCAGAGCCTGTTCCCTTCATCGAGAAGCTGGCGGCCTCAGTACATGCACTCCA  
TCAGGCCCAGGAGGACGAGCTACAGAAGCTGACCCAGCTCCGGGACTCCCTCCGAG  
GGACACTGCAGCTTGAGAGCAGAGAGGAACACCTGAGCCGGAAGAACTCAGGATGTG  
GCTATAGCATCCACCAGCACCAAGGCAACAAGCAGTTTGGGACGGAGAAAGTGGGCT  
TTCTATACAAGAAAAGTGACGGAATTGGAAGAGTCTGGCAGAAAAGGAAGTGTGGAGT  
CAAGTATGGCTGCCTGACCATCTCACACAGCACGATAAACCGGCCCGGTGAAGCTG  
ACCCTGCTGACGTGCCAAGTGAGGCCAAACCTGAGGAGAAAAAGTGTTCGACCTG  
GTGACCCACAACCGGACGTACCACTTTCAGGCAGAGGACGAGCACGAGTGTGAGGC  
GTGGGTGTCAGTGTTGCAGAACAGCAAGGACGAAGCCCTGAGCAGCGCCTTCCTCG  
GGGAGCCCAGCGCTGGCCCGGGGTCTGGGGGTCCGCCGGCCATGATGGGGAGCC  
GCACGACCTCACAAAGCTGCTCATCGCGGAGGTGAAGAGCAGGCCTGGGAATAGCCA  
GTGCTGCGACTCGGGGCTGCAGACCCACGTGGCTCAGCACCAACCTGGGCGTGC  
TCACCTGCATCCAGTGCTCGGGCGTCCACCGCAACTGGGCGTGCGCTTTTCGCGCA  
TGCACTCACTCACCTTGGACCTGCTGGGCCCTCCGAGTTGTTGCTGGCCTTGAACAT  
GGGAAACACGAGCTTCAATGAGGTCATGGAGGCCAGCTACCCTCACACGGCGGCCCC  
TAAACCTCAGCTGAGAGTGACATGGGCACCCGCAGGGACTACATTATGGCCAAGTAT  
GTGGAGCATAGGTTTGCACGCCGTGCACACCTGAGCCTCAGCGACTCTGGACAGCC  
ATTTGCAACAGGGACCTCCTGTCCGTACTGGAGGCCTTTGCCAATGGGCAGGACTTTG  
GACAGCCGCTGCCAGGGCCTGATGCACAGGCACCTGAAGAACTCGTCTTGCAATTGG  
CTGTCAAAGTCGCCAACCAAGGCTTCCCTGCTGCTGGTGGATTTCATCATCCAAACGG  
TGGTCACCTGGATGCCAAGGCTGCTGACGGGAACACGGCTCTGCACTACGCAGCACT  
CTACAACCAGCCCGACTGCCTCAAGCTGCTGCTGAAGGGGAGAGCTTTGGTTGGCAC  
AGTAAATGAAGCAGGCGAGACAGCTCTGGACATAGCCAGGAAGAAGCACCACAAGGA  
GTGTGAGGAGCTGCTGGAGCAGGCCAGGCGGGGACCTTTGCCTTCCCTCTACATGT  
GGACTACTCCTGGGTAATTTCCACAGAGCCTGGCTCTGACAGTGAGGAGGATGAGGAA  
GAGAAGCGCTGCTTGTGAAGCTCCCGGCCAGGCTCACTGGGCCAGTGGGAGGCT  
GGACATCAGCAACAAGACCTATGAGACTGTCCGAGCCTGGGAGCAGCCACCCCTCA  
GGCGGAGAGTGAGGACTGTCCCCGCCCTTGCCAGTCAAAAACCTCTTCTCGGACTTT  
GGTCCAAGGGTGTGCAAGACATGCCAGTGGAGATCGTTCTGAAGTCTCCAGCCTGAG  
TTCAGAGGCCCTGAGACCCCTGAGAGCCTGGGCAGTCCAGCCTCCTCCTCAGTCT  
GATGAGCCCCTTGGAACCTGGGGATCCAGCCAAGCCCCACCCAACCTCTGAAGAGGG  
CCTCCGAGAGCCCCCAGGCACCTCCAGACCCAGCCTGACATCCGGGACCACCCCTTC  
GGAGATGTACCTCCCCGTGAGATTCAGCTCCGAGAGCACTCGCTCCTATCGGCGGGG  
GGCGCGGAGCCCTGAAGATGGTCCCTCAGCCAGGCAGCCTCTGCCAGAAGGAACG  
TGCCGGTTGGCACTGAAGGAGATGGTCAAGGACTGGGAGTCTCCAGCAAGT  
CTGTGCAACTTTTGCAAGACTAGCTCTTGTGCTGGCCCCACATGCCCCATGCTAGGCC  
CCAATGTTGAGAGCTGGGACTTGAGCTCACAAAACCTGGGGAGCTGAGACATTTGTTCT  
CTTGGATCTCACTCTCTGTCCCTTGTGCCTCTGTAGCTGGCCTTCTCCTGCCACAG  
GCCATGCCTCTACCAAGGACACATGGCCTTCCCTGTTAGGGCTGATGGCGGTTCTTT

CCTATCTCATTACCCGCTAGGGGCCTGGGAGCCCTGTGGCTGGATCTGAGTGCTCCTG  
 AGCTGGCTTCAGCTGCAGAACTCTCAGTCCCTCATCAGATCGAGACTCTATTTCCCCC  
 GTCAGTCTGGGGGCTTCACAAGGGCAGGAGCCCTCCATCACTGACTTCCAGATCA  
 GGGACCCTGCCAAGTAGGGACTGTCTTCTCAGCCAGCCATTATTAGTCTAATATTCCT  
 TCACTAAATTCCAACCTCTATGTCTGGACCTGTGTTAGGCACTTCAGATACCACACGAGTA  
 AGACAAGGGCCCTGCAGGGGTGGTCTTTGGTGGAAAGCTGGTCTTAAGGGTTGGGC  
 TTGGGAATAGGCAGGGTCAGATTCCAGGGCATGGCTCTGGACTCAGCTGGTTTATACC  
 TATATGACCATTACAGTTGTCTACAGATCACATCCATTCTGGCTGGTCAACATGCATGCT  
 GTACTGGCTGTTAAATAAAAATATTCTGAATGTCACCTCTTTTGAAGGACAGCACAGCCT  
 TCCCTAGGCATTCTCCTATATTTCCCCAGCCAAATTGTAGAGTCAGATGCACCCACATTTG  
 CCTGTGTCCTTGATTTAGCAGGAAGGAAAGGAATAGTCGGGGTTGATGGATGCCCACT  
 TCTCTTCTCTTCTCTTGGTCAACTCAGGAGCCTTTTAGTCTGAGGGAATGGAGAGGCA  
 AAGAAAGAAGGGAGAGTAATAGAATTGGGAGGGCAGAGACTTAAGGGTTCTGCTTCCC  
 AGCCCTAGAAATTCTATCATTGCTCAGCCCCAATGAGAAAGCAGATACACCTAAGCCAT  
 CATCAACCACTAACATCTCAACTTGCCAGTTGCTGGGTGCTGGGCCCTGGCAGGAATG  
 GGCCAAGCCAAGCAGGGGAGACTAGAGAGCACCAATGGCCAACACAGCTGCCTGGC  
 TGGGGAGGCTGTGCTGTTTCCCTGGAGACCTGACTGGTCTGTGGTTCCACAGGAA  
 CAGGGTTGTCTTTGAGCCCCCAGTGCTGGTTTCATTCTCAGACTTGTTATTTCA  
 CTCATCTCTAATAAAGGATTGGGGGGTCACTT

## SEQ ID 2

DDEF1(amino acid)

MPEQFSVAEFLAVTAEDLSSPAGAAFAAKMPRYRGAALAREEILEGDQAILQRIKKAVRAI  
 HSSGLGHVNEEQYREAVESLGNHLSQNSHELSTGFLNLAVFTREVAALFKNLIQNLNNI  
 VSFPLDSLMMKGQLRDGRQDSKKQLEKAWKDYEAQMAKLEKERDRARVTGGIPGEVAQD  
 MQRERRIFQLHMCEYLLKAGESQMKQGPDLFQSLIKFFHAQHNFQDQGWKAAQSLFPFIE  
 KLAASVHALHQAQEDQLKLTQLRDSLRLGTLQLESREEHLRKNNSGCGYSIHQHQGNKQF  
 GTEKVGFLYKKSDGIRRVWQKRKCGVKYGLTISHSTINRPPVKLTLLTCQVRPNPEEKKC  
 FDLVTHNRITYHFQAEDEHECEAWWSVLQNSKDEALSSAFLGEPGAGPSWGSAGHDGE  
 PHDLTKLLIAEVKSRPGNSQCCDCGAADPTWLSTNLGVLTICQCSGVHRELGVRFSRMQS  
 LTLDLLGPSELLALNMGNSTFNEVMEALPSSHGGPKPSAESDMGTRRDYIMAKYVEHRF  
 ARRCTPEPQRLWTAICNRDLLSVLEAFANGQDFGQPLPGPDAQAPEELVHLAVKVANQA  
 SLPLVDFFIQQNGHLDKAADGNTALHYAALYNQPDCLKLLKGRALVGTVNEAGETALDIA  
 RKKHHKECEELLEQAQAGTFAPPLHVDYSWWISTEPGSDSEEDDEEEKRCLLKLPAQAHWA  
 SGRDLISNKTYETVASLGAATPQGESEDCPPPLPVKNSSRTLQVQGCARHASGDRSEVSSL  
 SSEAPETPESLGSPASSSSLMSPLEPGDPSQAPPNSEEGLREPPGTSRPSLTSGTTPSEM  
 YLPVRFSSSESTRSYRRGARSPEGDGPSARQPLPRRNVPGITEGDGSRTGSLPASSVQLLD

## SEQ ID 3

ZNFN3A1(nucleotide)

GTGCGCGCAGGGCGCAGGCGCGCGGGTCCCGGCAGCCCGTGAGACGCCCGCTGCT  
 GGACGCGGGTAGCCGTCTGAGGTGCCGGAGCTGCGGGAGGATGGAGCCGCTGAAG  
 GTGGAAAAGTTGCAACCGCCAACAGGGGAAACGGGCTGCGCGCCGTGACCCCGCT  
 GCGCCCCGGAGAGCTACTCTTCCGCTCGGATCCCTTGGCGTACACGGTGTGCAAGGG  
 GAGTCGTGGCGTCGTCTGCGACCGCTGCCTTCTCGGGAAGGAAAAGCTGATGCGATG  
 CTCTCAGTGCCGCGTCGCCAAATACTGTAGTGCTAAGTGTGAGAAAAAGCTTGGCCA  
 GACCACAAGCGGGAATGCAAATGCCTTAAAAGCTGCAAACCCAGATATCCTCCAGACT  
 CCGTTCGACTTCTTGGCAGAGTTGTCTTCAAACCTTATGGATGGAGCACCTTCAGAATCA  
 GAGAAGCTTTACTCATTTTATGATCTGGAGTCAAATATTAACAACTGACTGAAGATAAG  
 AAAGAGGGCCTCAGGCAACTCGTAATGACATTTCAACATTTTCATGAGAGAAGAAATACA  
 GGATGCCCTCTCAGCTGCCACCTGCCTTTTACCTTTTGAAGCCTTTGCAAAAGTATCT  
 GCAACTCTTCCACCATCTGTAATGCGGAGATGCAGGAAGTTGGTGTGGCCTATATCCC  
 AGTATCTCTTTGCTCAATCACAGCTGTGACCCCACTGTTGATTGTGTTCAATGGGCC  
 CCACCTCTTACTGCGAGCAGTCCGAGACATCGAGGTGGGAGAGGAGCTCACCATCTG  
 CTACCTGGATATGCTGATGACCAGTGAGGAGCGCCGGAAGCAGCTGAGGGACCAGTA

CTGCTTTGAATGTGACTGTTTCCGTTGCCAAACCCAGGACAAGGATGCTGATATGCTAA  
CTGGTGATGAGCAAGTATGGAAGGAAGTTCAGAATCCCTGAAAAAATTGAAGAACTG  
AAGGCACACTGGAAGTGGGAGCAGGTTCTGGCCATGTGCCAGGCGATCATAAGCAGC  
AATTCTGAACGGCTTCCCGATATCAACATCTACCAGCTGAAGGTGCTCGACTGCCCAT  
GGATGCCCTGCATCAACCTCGGCCTGTTGGAGGAAGCCTTGTTCTATGGTACTCGGACC  
ATGGAGCCATACAGGATTTTTTTCCAGGAAGCCATCCCGTCAGAGGGGTTCAAGTGA  
TGAAAGTTGGCAAACTGCAGCTACATCAAGGCATGTTTCCCCAAGCAATGAAGAATCTG  
AGACTGGCTTTTGATATTATGAGAGTGACACATGGCAGAGAACACAGCCTGATTGAAGA  
TTTGATTCTACTTTTAGAAGAATGCGACGCCAACATCAGAGCATCCTAAGGGAACGCAG  
TCAGAGGGAAATACGGCGTGTGTCTTTGTTGAATGCCTTATTGAGGTCACACACTCTAT  
GCTTTGTTAGCTGTGTGAACCTCTCTATTGGAATTCTGTTCCGTGTTTGTGTAGGTAA  
ATAAAGGCAGACATGTTTGCAAACCACAAGAATCATTAGTTGTAGAGAAGCACGATTAT  
AATAAATTCAAAACATTTGTTGAGGATGCCAAAAAAAAAAAAAAAAAAAAA

## SEQ ID 4

ZNFN3A1(amino acid)

MEPLKVEKFATANRGNGLRAVTPLRPGELLFRSDPLAYTVCKGSRGVVCDRCLLGKEKLM  
RCSQCRVAKYCSAKCQKKAWPDHKRECKCLKSCKPRYPDSVRLLGRVVFKLMDGAPS  
ESEKLYSFYDLESNINKLTEDKKEGLRQLVMTFQHFMRREEIQDASQLPPAFDLFEAFKVIC  
NSFTICNAEMQEVGVGLYPSISLLNHSCDPNCSIVFNPHLLRAVRDIEVGEELTICYLDM  
MTSEERRKQLRDQYCFECDFRCQTQDKDADMLTGDEQVWKEVQESLKKIEELKAHWK  
WEQVLAMCQAISSNSERLPDINIYQLKVLDCAMDACINLGLLEEALFYGTRTMEPYRIFFP  
GSHPVRGVQVMKVGKLQLHQGMFPQAMKNLRLAFDIMRVTHGREHSLIEDLILLLEECDA  
NIRAS

## SEQ ID 5

VANGL1

CTCGCTCACAAAAAATTGAGCCGGCCCTGGAGGCCTGGGGGGCGAGTCCGGTTGCG  
CCTCGGAGAGCGCAACAGGCAGAATTTGTTCTGTTGAAGAGTGGCTCCTCTTCTAAT  
TTCCAGACTCCTTGAGGTTTTAGGAGTCTGGTAGGTGAAATTTCTACCTCTAAGGAGA  
AACAGTACCTGCTCCTTCTCAAGCGCAAGCCCTCCATTGCTATGGATACCGAATCCAC  
TTATTCTGGATATTCTTACTATTCAAGTCATTGCAAAAAATCTCACAGACAAGGGGAAAG  
AACTAGAGAGAGACACAAGTCACCCCGGAATAAAGACGGCAGAGGGTCAGAAAAGTC  
TGTCACCATTC AACCTCCCCTGGAGAGCCCCTGTTGGGAAATGATTCTACTCGGACA  
GAGGAAGTTCAGGATGACAACCTGGGGAGAGACCACCGCCATCACAGGCACCTCG  
GAGCAGCATATCCCAAGAGGACATTGCCAGGATCAGCAAGGACATGGAGGACAGC  
GTGGGGCTGGATTGCAAACGCTACCTGGGCCTCACCGTCGCCTCTTTTCTTGGACTTC  
TAGTTTTCTCACCCCTATTGCCTTCATCCTTTTACCTCCGATCCTGTGGAGGGATGAG  
CTGGAGCCTTGTTGGCACAATTTGTGAGGGGCTCTTTATCTCCATGGCATTCAAACCTCT  
CATTCTGCTCATAGGGACCTGGGCACTTTTTTTCCGCAAGCGGAGAGCTGACATGCCA  
CGGGTGTTTGTGTTTCGTGCCCTTTTGTGGTCTCATCTTTCTCTTTGTGGTTTCTAT  
TGGCTTTTTTACGGGGTCCGCATTTTGGACTCTCGGGACCGGAATTACCAGGGCATTG  
TGCAATATGCAGTCTCCCTTGTTGGATGCCCTCCTCTTCATCCATTACCTGGCCATCGTC  
CTGCTGGAGCTCAGGCAGCTGCAGCCCATTGTCAGCTGCAGGTGGTCCGCTCCACC  
GATGGCGAGTCCCGCTTCTACAGCCTGGGACACCTGAGTATCCAGCGAGCAGCATTG  
GTGGTCTTAGAAAATTACTACAAAGATTTACCATCTATAACCCAAACCTCCTAACAGCC  
TCCAAATTCCGAGCAGCCAAGCATATGGCCGGGCTGAAAGTCTACAATGTAGATGGCC  
CCAGTAACAATGCCACTGGCCAGTCCCGGGCCATGATTGCTGCAGCTGCTCGGCGCA  
GGGACTCAAGCCACAACGAGTTGTATTATGAAGAGGCCGAACATGAACGGCGAGTAA  
GAAGCGGAAAGCAAGGCTGGTGGTTGCAGTGGAAGAGGCCTTCATCCACATTACGCG  
TCTCCAGGCTGAGGAGCAGCAGAAAGCCCCAGGGGAGGTGATGGACCCTAGGGAGG  
CCGCCAGGCCAATTTCCCTCCATGGCCAGGGCTCTCCAGAAGTACCTGCGCATCAC  
CCGGCAGCAGAACTACCACAGCATGGAGAGCATCCTGCAGCACCTGGCCTTCTGCAT  
CACCAACGGCATGACCCCCAAGGCCTTCTAGAACGGTACCTCAGTGCGGGGCCCCAC  
CCTGCAATATGACAAGGACCGCTGGCTCTCTACACAGTGGAGGCTTGTCAGTGATGAG  
GCTGTGACTAATGGATTACGGGATGGAATTGTGTTCTGCTTAAGTGCTTGGACTTCAG

CCTCGTAGTCAATGTGAAGAAAATCCATTCATCATACTCTCTGAAGAGTTCATAGACCC  
CAAATCTCACAAATTTGTCCTTCGCTTACAGTCTGAGACATCCGTTTAAAAGTTCTATATT  
TGTGGCTTTATTAATAAAAAAAGAAAAATATATAGAGAGATATGCAAAAAAATAAAAGAC  
AAAAACAAAAA

## SEQ ID 6

VANG1(amino acid)

MDTESTYSGYSYSSSHSKSHRQGERTRERHKSPRNKDGRGSEKSVTIQPPTGEPLGN  
DSTRTEEVQDDNWGETTTAITGTSEHSISQEDIARISKDMEDSVGLDCKRYLGLTVASFLGL  
LVFLTPIAFILLPPILWRDELEPCGTICEGLFISMAFKLLILLIGTWALFFRKRRADMPRVFVFR  
ALLLVLIFLVVSYWLFYGVRIILDSRDRNYQGIVQYAVSLVDALLFIHYLAIVLLELRQLQPMF  
TLQVVRSTDGESRFYSLGHLISQRAALVLENYYKDFIYNPNLLTASKFRAAKHMAGLKVY  
NVDGPSNNATGSRAMIAAAARRRDSSHNELYYEEAEHERRVKKRKARLVVAVEEAFIHIQ  
RLQAEQKQAPGEVMDPREAAQAIFPSMARALQKYLRIIRQQNYHSMESILQHLCITN  
GMTPKAFLERYLSAGPTLQYDKDRWLSTQWRLVSDEAVTNGLRDGIIVFLKCLDFSLVFN  
VKKIPFIILSEEFIDPKSHKFLRLQSETSV

## SEQ ID 7

LGN(nucleotide)

GGCACGAGGAAGAATCAGGAGCTTAGGATGTATTAACACCAACTCATTAATATACTAACC  
GGACAATGTTCTACAAACAATTCTACATTGTAAAGGACTGGATTGGCACAAAATAAAATA  
ATTTTATTTTATTCAGCTTATAATGACTCGATGGAGGAAAATTTGATAAGCATGAGAGA  
AGACCATTCTTTTCATGTTGTTACAGAATGGAAGCTTCTTGCTAGAGCTGGCCTTGG  
AAGGGGAACGTCTATGTAAATCAGGAGACTGCCGCGCTGGCGTGTCATTCTTTGAAGC  
TGCAGTTCAAGTTGGAAGCTGAAGACCTAAAAACACTTAGCGCTATTTACAGCCAGTTGG  
GCAATGCTTATTTCTATTTGCATGATTATGCCAAAGCATTAGAATATCACCATCATGATTTA  
ACCCTTGCAAGGACTATTGGAGACCAGCTGGGGGAAGCGAAAGCTAGTGGTAATCTG  
GGAAACACCTTAAAGTTCTTGGAATTTTGACGAAGCCATAGTTTGTGTGAGCGACA  
CCTAGATATTTCCAGAGAGCTTAATGACAAGGTGGGAGAAGCAAGAGCACTTTACAATC  
TTGGGAATGTGTATCATGCCAAAGGGAAGTTTGGTTGCCCTGGTCCCCAGGATGT  
AGGAGAATTTCCAGAAGAAGTGAGAGATGCTCTGCAGGCAGCCGTGGATTTTATGAG  
GAAAACCTATCATTAGTGACTGCTTTGGGTGACCGAGCGGCACAAGGACGTGCCTTTG  
GAAATCTTGGAACACACATTACCTCCTTGGAACCTCAGGGATGCAGTTATAGCTCAT  
GAGCAGCGTCTCCTTATTGCAAAGAATTTGGAGATAAAGCAGCTGAAAGAAGAGCATA  
TAGCAACCTTGGAATGCATATATTTCTTGGTGAATTTGAACTGCCTCGGAATACTAC  
AAGAAGACACTACTGTTGGCCCGACAGCTTAAAGACCGAGCTGTAGAAGCACAGTCTT  
GTTACAGTCTTGGAATACATATACTTTACTTCAAGACTATGAAAAGGCCATTGATTATCA  
TCTGAAGCACTTAGCAATTGCTCAAGAGCTGAATGATAGAATTGGTGAAGGAAGAGCAT  
GTTGGAGCTTAGGAAATGCATACACAGCACTAGGAAATCATGATCAAGCAATGCATTTT  
GCTGAAAAGCACTTGGAATTTCAAGAGAGGTTGGGGATAAAAGTGGTGAACCTAACAG  
CACGACTTAATCTCTCAGACCTTCAAATGGTTCTTGGTCTGAGCTACAGCACAAATAAC  
TCCATAATGTCTGAAAATACTGAAATTGATAGCAGTTTGAATGGTGTACTCCCCAAGTTG  
GGACGCCCGCATAGTATGGAAAATATGGAACCTTATGAAGTTAACACCAGAAAAGGTACA  
GAACCTGGAACAGTGAAATCTTGCTAAGCAAAAACCTCTTATTGCCAAACCTTCTGCAA  
AGCTACTCTTTGTCAACAGACTGAAGGGGAAAAAATACAAAACGAATTCCTCCACTAAA  
GTTCTCCAAGATGCCAGTAATTTCTATTGACCACCGAATTCCAAATTCCTCAGAGGAAAATC  
AGTGCAGATACTATTGGAGATGAAGGGTTCTTTGACTTATTAAGCCGATTTCAAAGCAAT  
AGGATGGATGATCAGAGATGTTGCTTACAAGAAAAGAACTGCCATACAGCTTCAACAAC  
AACTTCTTCCACTCCCCCTAAATGATGCTAAAAACATCATCTGTTCTGTGGTATCCCC  
CAACACGGATGAGTTTTAGATCTTCTTGCCAGCTCACAGAGTCGCCGTCTGGATGAC  
CAGAGGGCTAGTTTCAGTAATTTGCCAGGGCTTCGTCTAACACAAAACAGCCAGTCGG  
TACTTAGCCACCTGATGACTAATGACAACAAAGAGGCTGATGAAGATTTCTTTGACATCC  
TTGTAAATGTCAAGGATCCAGATTAGATGATCAAGATGTGCTCCACCACCTGCTACC  
ACAAAGGGTCCGACAGTACCAGATGAAGACTTTTTTCAGCCTTATTTACGGTCCCAGGG  
AAAGAGAATGGATGAACAGAGAGTTCTTTTACAAAGAGATCAAAACAGAGACACTGACT  
TTGGGCTAAAGGACTTTTTGCAAATAATGCTTTGTTGGAGTTTAAAAATTCAGGGAAAA

AATCGGCAGACCATTAGTTACTATGGATTTATTTTTTTCCTTTCAAACACGGTAAGGAAA  
CAATCTATTACTTTTTTCTTAAAAGGAGAATTTATAGCACTGTAATACAGCTTAAAATATT  
TTTAGAATGATGTAAATAGTTAA

SEQ ID 8

LGN(amino acid)

MREDHSFHVRYRMEASCLELALEGERLCKSGDCRAGVSFFEAQVVGTEDLKTLIAIYSQ  
LGNAYFYLHDYAKALEYHHDLTLARTIGDQLGEAKASGNLGNLTKVLGNFDEAIVCCQRH  
LDISRELNDKVGEARALYNLGNVYHAKGKSFGCPGPQDVGEFPPEVRDALQAAVDFYEEN  
LSLVTALGDRAAQGRAFGNLGNTHYLLGNFRDAVIAHEQRLLIAKEFGDKAAERRAYSNLG  
NAYIFLGEFETASEYYKKTLLARQLKDRAVEAQSCYSLGNTYTLLQDYEKADYHLKHLAIA  
QELNDRIGEGRACWSLGNAYTALGNHDQAMHFAEKHLEISREVGDKSGELTARLNLSDLQ  
MVLGLSYSTNNSIMSENTEIDSSLNGVLPKLGRRHSMENMELMKLTPEKVQNWNSILAK  
QKPLIAKPSAKLLFVNRLKGKKYKTNSSSTKVLQDASNSIDHRIPNSQRKISADTIGDEGFFDL  
LSRFQSNRMDDQRCCLQEKNCHTASTTSSSTPPKMMLKTSSVPVVSPTNDEFLLDASSQ  
SRRLLDDQRASFSNLPGLRLTQNSQSVLSHLMTNDNKEADEDFDILVKCQGSRLDDQRC  
PPPATTKGPTVPDEDFSLILRSQGKRMDEQRVLLQRDQNRDTDFGLKDFLQNNALLEFK  
NSGKKSADH

## CLAIMS

1. A marker for diagnosing cancer, wherein said marker is an expression product of any one of (i) the DDEFL1 gene set forth in SEQ ID NO: 1, (ii) the ZNFN3A1 gene set forth in SEQ ID NO: 3, (iii) the VANGL1 gene set forth in SEQ ID NO: 5, or (iv) the LGN gene set forth in SEQ ID NO: 7.
2. The marker according to claim 1, wherein a transcription product of the DDEFL1 gene is the protein set forth in SEQ ID NO: 2.
3. The marker according to claim 1, wherein a transcription product of the ZNFN3A1 gene is the protein set forth in SEQ ID NO: 4.
4. The marker according to claim 1, wherein a transcription product of the VANGL1 gene is the protein set forth in SEQ ID NO: 6.
5. The marker according to claim 1, wherein a transcription product of the LGN gene is the protein set forth in SEQ ID NO: 8.
6. A tool for detecting the marker according to claim 1, wherein said marker comprises (a) a nucleotide (i) complementary to the nucleotide sequence set forth in SEQ ID NO: 1, or to the complementary strand thereof, and, (ii) comprising at least 15 nucleotides, or (b) an antibody against the protein set forth in SEQ ID NO: 2.
7. A tool for detecting the marker according to claim 1, wherein said marker comprises (a) a nucleotide (i) complementary to the nucleotide sequence set forth in SEQ ID NO: 3, or to the complementary strand thereof, and, (ii) comprising at least 15 nucleotides, or (b) an


- antibody against the protein set forth in SEQ ID NO: 4.
8. A tool for detecting the marker according to claim 1, wherein said marker comprises (a) a nucleotide (i) complementary to the nucleotide sequence set forth in SEQ ID NO: 5, or to the complementary strand thereof, and, (ii) comprising at least 15 nucleotides, or (b) an antibody against the protein set forth in SEQ ID NO: 6.
  9. A tool for detecting the marker according to claim 1, wherein said marker comprises (a) a nucleotide (i) complementary to the nucleotide sequence set forth in SEQ ID NO: 7, or to the complementary strand thereof, and, (ii) comprising at least 15 nucleotides, or (b) an antibody against the protein set forth in SEQ ID NO: 8.
  10. A method for diagnosing cancer, comprising the step of measuring the expression level of any one of (i) the DDEFL1 gene set forth in SEQ ID NO: 1, (ii) the ZNFN3A1 gene set forth in SEQ ID NO: 3, (iii) the VANGL1 gene set forth in SEQ ID NO: 5, or (iv) the LGN gene set forth in SEQ ID NO: 7.
  11. A use of the DDEFL1 gene set forth in SEQ ID NO: 1, or the DDEFL1 protein set forth in SEQ ID NO: 2, as a target for developing drugs for treating cancer.
  12. A use of the ZNFN3A1 gene set forth in SEQ ID NO: 3, or the ZNFN3A1 protein set forth in SEQ ID NO: 4, as a target for developing drugs for treating cancer.
  13. A use of the VANGL1 gene set forth in SEQ ID NO: 5, or the VANGL1 protein set forth in SEQ ID NO: 6, as a target for developing drugs for treating cancer.
  14. A use of the LGN gene set forth in SEQ ID NO: 7, or the LGN protein set forth in SEQ ID NO: 8, as a target for developing drugs for treating cancer.
  15. A DNA comprising the nucleotide sequence set forth in




SEQ ID NO: 1.

16. A protein comprising the amino acid sequence set forth in SEQ ID NO: 2.
17. A DNA comprising the nucleotide sequence set forth in SEQ ID NO: 3.
18. A protein comprising the amino acid sequence set forth in SEQ ID NO: 4.


**Figure 1**



**Figure 2**




**Figure 3**




# Figure 4


a)



b)



c)



d)




Figure 5






Figure 6


a



b




**Figure 7**






**Figure 8**




**Figure 9**


**a**




**b**



**c**



**d**



**Figure 10**






Figure 11.


a




b




**Figure 12.**




**Figure 13.**



**Figure 14.**




**Figure 15**








**Figure 16**



**Figure 17**




**Figure 18**




**Figure 19**

**a**



**b**



**This Page is Inserted by IFW Indexing and Scanning  
Operations and is not part of the Official Record**

**BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☐ FADED TEXT OR DRAWING
- ☒ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☒ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: \_\_\_\_\_

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.**